

Research Article

Abundance and distribution of lantern fishes (Myctophiformes: Myctophidae) around San Pedro Martir Island, Gulf of California, during 2008

Mariana Díaz Santana-Iturríos¹, Deivis S. Palacios-Salgado² & César A. Salinas-Zavala³

¹Centro Interdisciplinario de Ciencias del Mar, Instituto Politécnico Nacional
Apartado Postal 592, La Paz, Baja California del Sur, México 23096

²Escuela Nacional de Ingeniería Pesquera, Colección Ictiológica (ENIP-UAN)
Apartado Postal 10, San Blas, Nayarit, México 63740

³Centro de Investigaciones Biológicas del Noroeste, Coordinación de Ecología Pesquera
P.O. Box 128, La Paz, Baja California del Sur, México 24090

ABSTRACT. Myctophids (Myctophidae) are a group of abundant mesopelagic fishes in the world's oceans and are known as the main feeding resource for several high trophic level predators. Changes in abundance may be related to population size of some commercially important species that feed on them. Only two of the myctophid species reported for the Gulf of California were found in the present study: *Benthosema panamense* and *Triphoturus mexicanus*. The highest abundance and biomass of myctophids were found during the warm season (June and September), with *B. panamense* being the most abundant species (20,954 ind 1000 m⁻³), as well as the one with highest biomass (17,165.8 g 1000 m⁻³). *B. panamense* had a size mode interval of 35-40 mm, while *T. mexicanus* presented a size mode interval of 40-45 mm; both species had negative allometric growth. During the temperate season (February and April) *B. panamense* was distributed in the northwest, west, and southern regions around the island, while *T. mexicanus* was found in the north, west, and southern regions. During the warm season *B. panamense* was found distributed around the entire island and *T. mexicanus* was found in the west, south, and east regions of the island. These species are common around San Pedro Martir Island, with the highest values of abundance and biomass occurring during summer upwelling's.

Keywords: abundance, *Benthosema panamense*, *Triphoturus mexicanus*, myctophids, Gulf of California.

Distribución y abundancia de peces linterna (Myctophiformes: Myctophidae) alrededor de isla San Pedro Mártir, golfo de California, durante el año 2008

RESUMEN. Los mictófidios (Myctophidae), son un grupo de peces mesopelágicos abundantes en los océanos del mundo, y se identifican como el principal recurso alimenticio de depredadores con alto nivel trófico. Los cambios en su abundancia pueden relacionarse con el tamaño poblacional de algunas especies de importancia comercial que depredan sobre ellos. En el presente trabajo, de las especies de mictófidios que se reportan dentro del golfo de California, sólo se encontraron dos: *Benthosema panamense* y *Triphoturus mexicanus*. En la temporada cálida (junio y septiembre) se presentaron las mayores abundancias y biomasa, siendo *B. panamense* la especie más abundante (20,954 ind 1000 m⁻³), así como también la de mayor biomasa (17,165.8 g 1000 m⁻³). *B. panamense* se encontró con una moda en el intervalo de tallas de 35-40 mm, *T. mexicanus* presentó una moda en el intervalo de tallas de 40-45 mm; ambas especies mostraron un tipo de crecimiento alométrico negativo. Con respecto a su distribución, durante la temporada templada (febrero y abril) *B. panamense* se distribuyó en la región noroeste, oeste y sur, mientras que *T. mexicanus* se encontró en las regiones norte, oeste y sur de la isla. Durante la temporada cálida *B. panamense* se encontró alrededor de toda la isla; *T. mexicanus* se encontró en las regiones oeste, sur y este de la isla. Estas especies son comunes alrededor de la isla San Pedro Mártir, con mayor abundancia y biomasa durante las surgencias de verano.

Palabras clave: abundancia, *Benthosema panamense*, *Triphoturus mexicanus*, mictófidios, golfo de California.

INTRODUCTION

Lantern fishes or myctophids (Myctophidae) are diverse and abundant in the world's oceans, represented by 240 species, grouped in 32 genera (Nelson, 2006). In the central-eastern Pacific Ocean region there are 20 genera and 45 species (Hart, 1973; Paxton *et al.*, 1995). They are generally small (2-30 cm), with large eyes, and have photophores in both the head and body with specific distribution patterns depending on the species (Frost & McCrone, 1979; Castellanos-Galindo *et al.*, 2006). These fishes are predominant in the mesopelagic zone, being one of the most abundant fish families in this zone; but can also be found over the continental shelf or around islands. Almost all of the species (except for the bathypelagic ones) migrate vertically from depths greater than 200 m (where they remain during the day) to the upper water column layer where they remain during the night (Paxton *et al.*, 1995). The highest abundance of almost all myctophid species occur between depths of 300 and 1200 m during the day and between 10 and 100 m during the night (Nelson, 2006).

It has been identified that myctophids are the main feeding resource for several high trophic level predators (Pakhomov *et al.*, 1996; Bañon *et al.*, 2000; Phillips *et al.*, 2001). In stomach contents of jumbo squid, *Dosidicus gigas* (D'Orbigny, 1835), in the Gulf of California, it has been found that its diet is mainly dominated by *Benthosema panamense* (Tåning, 1932) and *Triphoturus mexicanus* (Gilbert, 1890) (Markaida & Sosa-Nishizaki, 2003; Markaida, 2006; Markaida *et al.*, 2008).

Jumbo squid is an important species in the food web of the pelagic ecosystem in the Gulf of California (Rosas-Luis *et al.*, 2008). In the past decade, this species has gained greater importance given that it represents one of the few fisheries whose abundance has surprisingly increased, although squid shows unpredictable density fluctuations depending on location (Martinez-Aguilar *et al.*, 2006).

Changes in abundance of prey likely influence distribution and population size of its predators as well as the energy-flow patterns and the performance of the pelagic ecosystems (Cury *et al.*, 2000). Knowing myctophid fish abundance variations is important given that they are an elementary component of the trophic web in the pelagic ecosystem of the central region of the Gulf of California, where currently, energy-flow seems to be regulated by an active myctophid predator, the jumbo squid. The present research was carried out to determine spatial and temporal distribution of myctophid fishes around San Pedro Martir Island in the Gulf of California, during

2008, as well as to determine myctophid species composition and to describe the size structure of these fishes in one of the main aggregation zones of the jumbo squid.

MATERIALS AND METHODS

San Pedro Martir Island is located in the central region of the Gulf of California at 28°23'N and 112°20'W (Fig. 1). Its total area is 2.8 km², and it is found equidistant to three regions with the highest primary productivity: summer/spring upwelling in the Guaymas area, winter/autumn upwelling outskirts in the Santa Rosalia area, and the annual surrounding tidal mix of Ballenas Channel. Given that San Pedro Martir Island is located near the temperate current dynamics of the Ballenas Channel area and near warmer waters located southwards of the island, hence affected by these water masses (Tershy *et al.*, 1992).

Biological material used in this survey comes from four oceanographic cruises focused in capturing juveniles and paralarvae of jumbo squid, *Dosidicus gigas* (DG) in the Gulf of California (GOLCA). Cruises were conducted in 2008 during February (DGGOLCA0208), April (DGGOLCA0408), June (DGGOLCA0608), and September (DGGOLCA0908), comprising two seasons: temperate (February and April) and warm (June and September), on board of a research vessel called "BIP XII", in the area of San Pedro Martir Island, Gulf of California. On each cruise eight stations were sampled, for a total of 32 stations (Table 1).

Myctophid samples were obtained using an Isaacs-Kidd midwater trawl with 3 m mouth width and 33 m length, procedure consisted of oblique trawling at an approximated speed of 1 knot. Samples were fixated in 10% formaldehyde. Samples were later identified using a stereoscopic microscope Zeiss Stemi DV4, and fishes from family Myctophidae were separated. Identification was made following Wisner (1976) and Paxton *et al.* (1995) proposed criteria. Standard length (SL) was taken for each specimen using an ichthyometer (1 mm). Total weight (TW) was also measured in grams using an analytical balance OHAUS EP114C (0.001 g). Weight and size frequency were taken per species and per cruise without separating males from females because of the absence of external characters that showed sexual dimorphism.

To define the relation of weight increase according to myctophid's length, the following linear equation was applied: $Y = a \cdot X^b$, where: Y = dependent variable (weight), a = proportionality coefficient, X = indepen-




Figure 1. Study area located around San Pedro Martir Island.

dent variable (length), and b = allometric coefficient (Sparre & Venema, 1997).

To evaluate growth type, a t-Student test was performed for each of the “ b ” values obtained from the model, and compared with $b = 3$, the theoretical value representative of isometric growth.

Relative abundance was estimated through the filtered volume method (FVM). The midwater trawl method covered a well defined water sector of the pelagic zone; the distance length of each sector was calculated by dividing velocity (knots) and trawling period. FVM consists in multiplying sector’s length by net’s mouth width and length.

Distribution maps were made with estimated relative abundance (RA) values, from the number of captured specimens, according to filtered water volume in terms of number of individuals 1000 m^3 , per analyzed cruise and per season (temperate and warm) using Surfer program (Golden Software V 9).

Relative biomass (RB) was estimated from weight of captured organisms per filtered water volume unity (g m^3). Biomass values were standardized to 1000 m^3 .

RESULTS

A total of 3,510 organisms were sampled from the four cruises made, taxonomic composition of family Myctophidae was found represented by two species: *Benthosema panamense* with 1606 organisms and *Triphoturus mexicanus* with 1904 organisms. The general size range for the species *B. panamense* was between 1-55 mm (mode = 35-40 mm) and for *T. mexicanus* the size interval was 10- 65 mm (mode = 40-45 mm) (Fig. 2).

Results of length-weight relationship analysis for each species showed that allometric coefficient obtained for *B. panamense* (2.09) and *T. mexicanus* (2.79), according to t-Student test, statistically deferred from the theoretical value of 3 ($P < 0.05$), therefore, growth in these species is considered to have negative allometry (Fig. 3).

According to the distribution and relative abundance (RA), during temperate season, *B. panamense* was distributed in regions of the northwest, west, and south of the island with total RA value of 338 ind 1000 m^3 . *T. mexicanus* was found in regions north

Table 1. Characteristics of the oceanographic cruises made around San Pedro Martir Island in the central part of the Gulf of California. A: *Bentosema panamense*, B: *Triphoturus mexicanus*.

Cruise	Station	Date	Time	Depth (m)	Latitude (N)	Longitude (W)	Biomass (g 1000 m ⁻³)		Abundance (ind 1000 m ⁻³)	
							A	B	A	B
DGGOLCA0208	1	08/02/2008	19:36	118	28° 27.296	112° 27.912	26.17	140.71	98.17	229.07
	2	08/02/2008	21:03	165	28° 22.828	112° 11.815	0	0	0	0
	3	08/02/2008	00:10	236	28° 16.410	112° 17.380	0	10.90	0	21.81
	4	09/02/2008	03:16	236	28° 22.023	112° 25.043	0	0	0	0
	5	09/02/2008	09:24	236	28° 22.523	112° 22.059	0	0	0	0
	6	09/02/2008	11:16	165	28° 26.335	112° 18.590	0	0	0	0
	7	09/02/2008	15:00	95	28° 22.242	112° 14.254	0	0	0	0
	8	09/02/2008	16:41	236	28° 19.394	112° 17.239	0	0	0	0
DGGOLCA0408	1	19/04/2008	13:37	118	28° 27.748	112° 17.110	0	0	0	0
	2	19/04/2008	15:38	212	28° 23.020	112° 11.645	0	0	0	0
	3	19/04/2008	18:20	212	28° 16.245	112° 17.229	0	0	0	0
	4	19/04/2008	20:46	212	28° 21.448	112° 29.662	47.99	1415.88	87.26	2443.44
	5	19/04/2008	22:40	212	28° 22.928	112° 22.023	17.45	167.98	21.81	218.16
	6	20/04/2008	00:30	118	28° 26.814	112° 20.120	80.72	305.43	87.26	283.61
	7	20/04/2008	02:40		28° 22.283	112° 15.033	0	0	0	0
	8	20/04/2008	04:20	212	28° 18.955	112° 17.737	34.90	1644.95	43.63	2334.35
DGGOLCA0608	1	25/06/2008	20:42	106	28° 27.961	112° 17.165	3506.00	0	4581.45	0
	2	25/06/2008	22:46	148	28° 23.133	112° 12.096	5030.67	263.284	6634.18	475.99
	3	26/06/2008	01:00	318	28° 16.624	112° 17.299	7.55	2976.26	8.39	5655.49
	4	26/06/2008	03:16	318	28° 21.887	112° 24.570	147.78	3271.40	263.90	6460.48
	5	26/06/2008	05:15	318	28° 22.767	112° 22.771	317.89	129.02	635.79	205.69
	6	26/06/2008	06:49	318	28° 26.110	112° 18.952	7649.38	0	7853.91	0
	7	26/06/2008	08:48	318	28° 22.226	112° 14.407	31.00	0	34.44	0
	8	26/06/2008	10:05	318	28° 19.507	112° 17.289	381.78	378.15	666.61	618.13
DGGOLCA0908	1	27/09/2008	08:55	148	28° 27.932	112° 17.126	0	0	0	0
	2	27/09/2008	11:45	212	28° 23.440	112° 11.762	0	0	0	0
	3	27/09/2008	14:39	255	28° 15.967	112° 17.038	0	0	0	0
	4	27/09/2008	18:33	255	28° 21.494	112° 25.855	41.81	93.35	218.16	418.14
	5	27/09/2008	20:37	255	28° 21.766	112° 22.983	21.17	1027.93	38.49	1366.73
	6	27/09/2008	23:16	148	28° 25.769	112° 20.645	0	0	0	0
	7	28/09/2008	02:10	85	28° 20.304	112° 14.828	0	46.74	0	93.49
	8	28/09/2008	03:49	255	28° 18.573	112° 14.891	30.67	2200.73	20.45	3170.20

west, and south of the island with total RA value of 5,530 ind 1000 m³, presenting higher RA in southern and western regions. During the warm season *B. panamense* was found all around the island with total RA value of 20,956 ind 1000 m⁻³, this being the highest RA value found in this study, while *T. mexicanus* was found in regions west, south, and east

of the island with total RA value of 18,464 ind 1000 m⁻³ (Fig. 4).

Highest relative biomass (RB) value for both species was found during the DGGOLCA0608 cruise being 17,072 g 1000 m⁻³ for *B. panamense* and 7,018.1 g 1000 m⁻³ for *T. mexicanus*. During the warm season, the highest RB was found for both species (*B*




Figure 2. a) Size frequency of *Benthosema panamense*, b) size frequency of *Triphoturus mexicanus*.

panamense: 17,165.8 g 1000 m⁻³ and *T. mexicanus*: 10,386.9 g 1000 m⁻³), with lower biomass in the temperate season with 207.3 g 1000 m⁻³ and 3,685.9 g 1000 m⁻³, respectively (Table 1).

DISCUSSION

Present study contributes to the understanding of some of the biological aspects of the most abundant myctophid species in the central region of the Gulf of California (Aceves-Medina *et al.*, 2003).

Of the 16 myctophid species present in the Gulf of California (Castro-Aguirre & Balart, 1996), two species were found in the cruises made during this study: *Benthosema panamense* and *Triphoturus mexicanus*. These species have been reported as abundant in fish larvae studies in the Gulf of California (Moser *et al.*, 1974; Avalos-Garcia *et al.*, 2003; Aceves-Medina *et al.*, 2004), also, both have been part of the major constituents of mid-water trawling expeditions made in this area, where the center of its distribution is in the central part of the gulf (Lavenberg & Fitch, 1966; Robison, 1972; Moser *et al.*, 1974), consistent to what was found in the present research. The low species richness of the collected specimens in this study may be due to the fact that the number of fish species living in the Gulf of California, decreases from south to north (Moser *et al.*, 1974). Or else, because of the employed sampling depths range (85-318 m), given that myctophids depth distribution in the water column varies widely according to each species and even within a species depending on its ontogenetic development (Paxton *et al.*, 1995). Another factor that may be influencing low species richness in the surveys of this study could be the lack of prey availability for other myctophid species feeding in sampled area (Frost & McCrone, 1979).

Data of size structure shows that *T. mexicanus* had an interval size mode of 40-45 mm for SL (40-45 mm SL), while *B. panamense* presented an interval size mode of 35-40 mm for SL (Fig. 2). It has been reported that there are ontogenetic changes in depth distribution patterns as well as daily vertical migration of myctophid fish, the trend is that organism's size increases according to depth (Frost & McCrone, 1979). Probably, myctophids with larger sizes were distributed at greater depths in the water column than




Figure 3. a) Standard length-weight relation of *Benthosema panamense*, b) standard length-weight relation of *Triphoturus mexicanus*.




Figure 4. Relative abundance and distribution per season for *Benthosema panamense* and *Triphoturus mexicanus*.

the depth range covered in this survey. Nevertheless, despite depths of myctophid fish sampling during the different cruises, size intervals were diverse (85-318 m) for both myctophid species and were relatively constant, this may be due to fishing gear selectivity.

Regarding the aforementioned, *B. panamense* is found between 200 and 300 m depth during the day and at nights it migrates to shallow waters above 100 m and *T. mexicanus* is found at the oxygen minimum layer during the day and at nights it rises in less aggregated shoals also above 100 m (Robison, 1972).

Energetic cost of swimming for most fishes decreases with size increase among and within species; such a relationship indicates that it is more expensive for a small fish to move 1 g of its corporal mass a given distance than it is for a larger fish (Helfman *et al.*, 2009), in this study, the allometric negative growth that both myctophid species showed may be due to the daily vertical migration that these fishes perform (Robison, 1972). They live in a viscous medium through which they move daily: myctophids in the upper 100 m depth at night are active and swim horizontally, also, these fishes are capable of rapid evasive movements, while those found at depths below 100 m are immobile and vertically oriented (Barham, 1971), perhaps the elongated body might be an advantage in terms of locomotion.

According to abundance and considering that vertical migration patterns of fishes from the Myctophidae family are species specific (Watanabe *et al.*, 1999), and that they are found at specific depths, variations in presence or absence of species in sampling stations, the capture of different sizes, as well as in abundance within a cruise, could possibly be due to a bias error for not having established a day/night schedule for sampling or considered species specific distribution depths. Doing so would have assured that all of the distributional range was covered when sampling.

One of the main natural fertilization mechanisms of the Gulf of California is the wind-induced upwelling (Alvarez-Borrego, 2010); winds that change directions seasonally and the strong tides (because of sea bottom shape) cause strong upwelling (Maluf, 1983); these upwelling during the winter (December-May) are caused by northern winds that drive the water surface layers out of the Gulf, having a significant effect on phytoplankton communities, this has been determined by the high chlorophyll concentration that has been found in these areas ($>100 \text{ mg m}^{-3}$) (Roden, 1958; Alvarez-Borrego, 2010). It was found in this study that during the temperate season myctophids were located towards the northwestern and southern regions of San Pedro Martir Island; also,

in this season the lowest abundance values for both species were obtained. Upwelling's during the summer (June-November), are caused by southern winds that drive oceanic water into the gulf at great depths, however these winds have a weak effect on phytoplankton communities and biomass because of the low chlorophyll concentration values ($\approx 0.5 \text{ mg m}^{-3}$) (Roden, 1958; Álvarez-Borrego, 2010), during the warm season myctophids were present all around the island and abundances were significantly higher. Myctophid abundances in this study might be determined by other biological factors rather than just feeding, it could possibly be related to reproduction, given that it has been found in previous studies, that these species show marked seasonal abundance and its larvae are found concentrated along the western side of the gulf, decreasing in density towards the east during the temperate season (Moser *et al.*, 1974). In the central portion of the Gulf of California, high concentration of sperm whales around San Pedro Martir Island likely indicates that there is probably a high concentration of jumbo squid, *Dosidicus gigas* (Gilly *et al.*, 2006), which feeds on myctophid fishes (Markaida & Sosa-Nishizaki, 2003).

The largest biomass values during the temperate season was determined to be *T. mexicanus* 3,686 g 1000 m³ and during the warm season *B. panamense* biomass was 17,166 g 1000 m³, the latter being the species having the largest biomass values from all cruises. Aforementioned explains why *D. gigas* diet in the Gulf of California is dominated firstly by *B. panamense* and secondly by *T. mexicanus* (Markaida & Sosa-Nishizaki, 2003; Markaida, 2006; Markaida *et al.*, 2008).

According to the analysis carried out it can be concluded that *B. panamense* and *T. mexicanus* are common around San Pedro Martir Island; both species presented negative allometric growth, indicating that length and width increase is not proportional; with highest abundance occurring for both species during the warm season. Of the two species, *B. panamense* was the most abundant.

ACKNOWLEDGEMENTS

We would like to thank CONACyT for funding "Proyecto Calamar" No. 165-C which permitted us to carry out the surveys. To Jasmin Granados-Amores, for providing invaluable help in conducting the study; to the captain and crew of "BIP XII", for assisting us with sampling procedures.

REFERENCES

- Aceves-Medina, G., S.P.A. Jimenez-Rosenberg, A. Hinojosa-Medina, R. Funes-Rodriguez, R.J. Saldierna, D. Lluch-Belda, P.E. Smith & W. Watson. 2003. Fish larvae from the Gulf of California. *Sci. Mar.*, 67(1): 1-11.
- Aceves-Medina, G., S.P.A. Jimenez-Rosenberg, A. Hinojosa-Medina, R. Funes-Rodriguez, R.J. Saldierna & P.E. Smith. 2004. Fish larvae assemblages in the Gulf of California. *J. Fish Biol.*, 65: 832-847.
- Alvarez-Borrego, S. 2010. Physical, chemical, and biological oceanography in the Gulf of California. In: R.C. Brusca (ed.). *The Gulf of California: biodiversity and conservation*. University Arizona Press, Arizona, pp. 24-48.
- Avalos-Garcia, C., L. Sanchez-Velasco & B. Shirasago. 2003. Larval fish assemblages in the Gulf of California and their relation to hydrographic variability (autumn 1997-summer 1998). *Bull. Mar. Sci.*, 72(1): 63-76.
- Bañon, R., S. Cerviño & J.M. Campelos. 2000. Composición, distribución y descripción de mictófidios (Pisces, Myctophidae) encontrados en Flemish Cap (Atlántico noroeste) en verano de 1998. *Bol. Inst. Esp. Oceanogr.*, 17(3-4): 287-294.
- Barham, E.G. 1971. Deep-sea fishes: lethargy and vertical orientation. In: G.B. Farquhar (ed.). *Proceedings of an international symposium on biological sound scattering in the ocean*. Maury Center for Ocean Science, Washington, Rept. Mc-005, pp. 100-118.
- Castellanos-Galindo, G.A., E.A. Rubio, B.S. Beltran-Leon & C.C. Baldwin. 2006. Check list of stomiiform, aulopiform and myctophiform fishes from Colombian waters of the Tropical Eastern Pacific. *Biota Col.*, 7(2): 245-262.
- Castro-Aguirre, J.L. & E.F. Balart. 1996. Contribución al conocimiento del origen y relaciones de la ictiofauna de aguas profundas del golfo de California, México. *Hidrobiologica*, 6(1-2): 67-76.
- Cury, P., A. Bakun, R.J.M. Crawford, A. Jarre, R.A. Quñones, L.J. Shannon & H.M. Verheye. 2000. Small pelagics in upwelling systems: patterns of interaction and structural changes in "wasp-waist" ecosystems. *ICES J. Mar. Sci.*, 57: 603-618.
- Frost, B.W. & L.E. McCrone. 1979. Vertical distribution, diel vertical migration and abundance of some mesopelagic fishes in the eastern subantarctic Pacific Ocean in summer. *Fish. Bull.*, 76(4): 751-770.
- Gilly, W.F., U. Markaida, C.H. Baxter, B.A. Block, A. Boustany, L. Zeidberg, K. Reisenbichler, B. Robinson, G. Bazzino & C. Salinas. 2006. Vertical and horizontal migrations by the jumbo squid

- Dosidicus gigas* revealed by electronic tagging. Mar. Ecol. Prog. Ser., 324: 1-17.
- Hart, J.L. 1973. Pacific fishes of Canada. Fish. Res. Bd. Can. Bull., 189: 740 pp.
- Helfman, G.S., B.B. Collette, D.E. Facey & B.W. Bowen. 2009. The diversity of fishes. Wiley-Blackwell, London, 720 pp.
- Lavenberg, R.J. & J.E. Fitch. 1966. Annotated list of fishes collected by midwater trawl in the Gulf of California, March-April 1964. Cal. Fish Game, 52(2): 92-110.
- Maluf, L.Y. 1983. Physical oceanography. In: T.J. Case & M.L. Cody (eds.). Island biogeography in the Sea of Cortez. University of California Press, California, 508 pp.
- Markaida, U. 2006. Food and feeding of jumbo squid *Dosidicus gigas* in the Gulf of California and adjacent waters after the 1997-1998 El Niño event. Fish. Res., 79: 16-27.
- Markaida, U. & O. Sosa-Nishizaki. 2003. Food and feeding habits of jumbo squid *Dosidicus gigas* (Cephalopoda: Ommastrephidae) from the Gulf of California, Mexico. J. Mar. Biol. Assoc. U.K., 83: 507-552.
- Markaida, U., W.F. Gilly, C.A. Salinas-Zavala, R. Rosas-Luis & J.A.T. Booth. 2008. Food and feeding of jumbo squid *Dosidicus gigas* in the central Gulf of California during 2005-2007. CalCOFI Rep., 49: 90-103.
- Martinez-Aguilar, S., J.G. Diaz-Uribe & M.O. Nevarez. 2006. Calamar gigante del Océano Pacífico. Sustentabilidad y Pesca Responsable en México, Evaluación y Manejo. INAPESCA, México, 544 pp.
- Moser, H.G., E.H. Ahlstrom, D. Kramer & E.G. Stevens. 1974. Distribution and abundance of fish eggs and larvae in the Gulf of California. CalCOFI Rep., 17: 112-128.
- Nelson, J.S. 2006. Fishes of the world. John Wiley & Sons, Otawa, 613 pp.
- Pakhomov, E.A., R. Perissinotto & C.D. McQuaid. 1996. Prey composition and daily rations of myctophid fishes in the Southern Ocean. Mar. Ecol. Prog. Ser., 134: 1-14.
- Paxton, J.R., R.J. Lavenberg & C. Sommer. 1995. Myctophidae: linternillas. In: W.F. Schneider, C. Sommer, K.E. Carpenter & V.H. Niem (eds.). Guía FAO para identificación de especies para los fines de la pesca. Pacífico Centro-Oriental. FAO, Rome, 3: 1315-1321.
- Phillips, K.L., G.D. Jackson & P.D. Nichols. 2001. Predation on myctophids by the squid *Moroteuthis ingens* around Macquarie and Heard Islands: stomach contents and fatty acids analyses. Mar. Ecol., 215: 179-189.
- Robison, B.H. 1972. Distribution of the midwater fishes of the Gulf of California. Copeia, 1972(3): 448-461.
- Roden, G.I. 1958. Oceanographic and meteorological aspects of the Gulf of California. Pac. Sci., 17: 21-45.
- Rosas-Luis, R., C.A. Salinas-Zavala, V. Koch, P. Del Monte-Luna & M.V. Morales-Zarate. 2008. Importance of jumbo squid *Dosidicus gigas* (D'Orbigny, 1835) in the pelagic ecosystem of the central Gulf of California. Ecol. Mod., 218: 149-161.
- Sparre, P. & S.C. Venema. 1997. Introducción a la evaluación de recursos pesqueros tropicales. FAO, Past 1, N° 306.1, Rev. 1, 420 pp.
- Tershy, B., D. Breese, A. Angeles, M. Cervantes, M. Mandujano, E. Hernandez & A. Cordoba. 1992. Natural history and management of Isla San Pedro Martir, Gulf of California. Reporte de Conservación Internacional, Sonora, 83 pp.
- Watanabe, H., M. Moku, K. Kawaguchi, K. Ishimaru & A. Ohno. 1999. Diel vertical migration of myctophid fishes (Family Myctophidae) in the transitional waters of the western North Pacific. Fish. Oceanogr., 8(2): 115-127.
- Wisner, R.L. 1976. The taxonomy and distribution of lanternfishes (Family Myctophidae) of the eastern Pacific Ocean. Naval Ocean Research and Development Activity, California, 229 pp.

Received: 13 July 2012; Accepted: 7 May 2013